

# Performance Equine Vets

3827 Charleston Hwy  
Aiken, SC 29801  
803.641.0644 888.874.7569  
www.PerformanceEquineVets.com



## **Frozen Equine Semen Handling & Thawing Instructions**

**This frozen semen has been shipped to you in a special Vapor “Dry” Shipper for Liquid Nitrogen and can only maintain proper storage temperature for a limited period of time. If you do not use the semen in this tank within a few days of arrival, PLEASE transfer it to your own liquid nitrogen storage tank to avoid damage to the semen. This container must be returned using the provided return label within 7 days, or a late fee of \$25.00 per day will apply to the mare owner.**

### **Special Precautions**


- Always keep this shipper in the upright position. Do Not splash or pour liquid nitrogen on your skin as it will burn.
- The transferring of straws to another liquid storage container should only be performed by a technician experienced in the handling of frozen semen. *Exposure of of the frozen straws for even a few seconds to room temperature may cause partial thawing and permanent damage to the semen.*
- Do Not put Goblet into water bath with straws still contained in it as they will thaw too slowly.
- Do Not pull one straw at a time out of goblet and plunge into water bath, as it will take too long.
- Do Not use less than one liter of water in 37°C bath. 1+ to 2 L of volume is a more stable temperature.
- Do Not pour liquid nitrogen into your bath as it will change your temperature by a few degrees quickly.

### *Materials needed for semen thawing:*

- *37°C constant temperature water bath (Do Not use water temperature that is above 37°C or below 35.5°C.)*
  - *Long handled tweezers or hemostats (pre-cool by holding down inside tank 6 to 10 inches for a few minutes prior to goblet removal.)*
  - *Sterile insemination supplies warmed to 37°C (use only ALL plastic syringes)*
  - *Protective eyewear and cyro-protective or leather gloves.*
1. Prepare mare aseptically prior to opening the container. Each dose is packaged in a goblet that is loaded on a cane.
  2. Lift tank canister with aluminum canes and /or goblet close to the top of the tank and remove the goblet. Do not lift the canister above the frost line at any time. (If the dose you need is the only one remaining on the cane, you may remove the entire cane). BE SURE to pre-cool your hemostats or long tweezers. Using only a few seconds: Remove goblet, pour out any liquid nitrogen, then quickly pull **All** straws out of goblet and drop or plunge straws into the 37°C bath for 30 Seconds. Use Cryogenic or leather gloves to handle goblet / cane. The steel ball sealed straws will bob in bath water with steel ball end down and critoseal straws may sink to bottom. Sink any floaters immediately.
  3. The individual straw may remain in the water bath while remaining straws are removed from the tank and thawed. Once all of the straws for the insemination dose have been thawed, remove them all together and wipe the straws completely with a paper towel in the palm of your hand. Keep straws in your warm hand for stable warm temperature. Water leaking into the straws is spermicidal. Do not lay straws on counter as it could be about 10+° cooler.
  4. Holding the straws vertically so that the air bubble is positioned on the top of the straw (the top is crimped or sealed with a sealing ball) cut off the crimped or sealed end with clean uncontaminated scissors. Next, invert all the straw open ends into a sterile, pre-warmed, 37°C glass or plastic container. Now cut off the larger cotton/pvc plug end of the straw at semen/cotton line, allowing semen to drain or flow into the warm tube.
  5. **We Do Not recommend extending semen as it may do damage to sperm.**

# Performance Equine Vets

3827 Charleston Hwy  
Aiken, SC 29801  
803.641.0644  
888.874.7569  
[www.PerformanceEquineVets.com](http://www.PerformanceEquineVets.com)



Stallion: \_\_\_\_\_

Straws per dose: \_\_\_\_\_

Number of doses sent: \_\_\_\_\_

## **Mare Management for Insemination with Frozen Semen**

Prior to insemination verify the mare is a suitable candidate for AI with frozen semen by performing a routine reproductive examination that may include culture, cytology, and biopsy when indicated. Data suggests that aged (>15 years) or repeatedly barren mares may have a reduced pregnancy rate and are not the best candidates for AI with frozen semen.

Once the mare comes into estrus, ultrasound daily to monitor follicular activity. Upon detection of a large (35-40mm) pre-ovulatory follicle, administer hCG or deslorelin or other induction agent of choice.

### **If only one dose is available for insemination:**

- Examine the mare via ultrasound at 6 hour intervals starting 12-24 hours after hCG/deslorelin/other administration.
- Inseminate the single dose of frozen semen as soon as ovulation is detected.
- It is extremely important that mares being inseminated post-ovulation are inseminated within 6 hours of ovulation. A significant reduction in fertility will occur if mares are inseminated more than 6 hours post ovulation.
- Semen may be deposited in uterine body or deep horn, gently plunge syringe to inseminate the entire amount of semen within minutes of loading pipette using 3cc of air. Hold pipette at a down angle while still in uterus if possible, which will allow the pipette wall coated semen (about one fourth of a straws volume) to drain down into the end of pipette. After a minute or two, plunge the remaining 2cc air and remaining drained semen. A few drops of semen will always be left in the AI pipette when held up right for a period after AI and can be examined if desired. Also, the unloaded empty straws can be taped end down on a warm glass slide after unloading straws to examine the unused small droplet with a cover slip. Thawed frozen semen microscopic evaluation may look best at about 5 minutes on a 37°C warmed glass slide.

\*\*\* In our hospital, mares are palpated every 2-4 hours once ovulation is pending, to more accurately time insemination and therefore, increase pregnancy rates.

\*\*\**All mares in our hospital receive a uterine flush and oxytocin administration ~6 hours post insemination.*

A general rule for mares inseminated with frozen semen is to inseminate within 12 hours prior to and/or within 6 hours after ovulation. This protocol ensures that viable sperm are in the oviduct during that interval for any mare ovulating within a period of 18 to 52 hours following administration of hCG or deslorelin or other induction agent.

**Feel free to call if you have any questions or concerns, we are more than happy to help guide you through the process.**

*~Thanks, Performance Equine Vets Staff*

**803.641.0644**

**888.874.7569**

**[info@PerformanceEquineVets.com](mailto:info@PerformanceEquineVets.com)**